

# How a leaf gets its shape

## Jihyun Moon and Sarah Hake

Leaves are formed from a group of initial cells within the meristem. One of the earliest markers of leaf initiation is the down-regulation of *KNOX* genes in initial cells. Polar auxin activity, MYB and LOB domain transcription factors function to keep *KNOX* out of the initiating leaf. If *KNOX* genes are expressed in initial cells, leaves fail to form. As the leaf grows away from the meristem, its shape is determined by growth in three axes, proximal–distal, abaxial–adaxial and medial–lateral. HD-ZIP III, KANADI and the small RNA pathway play a significant role in the latter two axes. *KNOX* proteins play a role in the proximal–distal axis. Although genetic networks are conserved between monocots and dicots, the outcome in leaf shape often differs.

### Address

Plant Gene Expression Center, USDA-ARS, University of California, Berkeley, 800 Buchanan St, Albany, CA 94710, United States

Corresponding author: Hake, Sarah ([hake@berkeley.edu](mailto:hake@berkeley.edu))

**Current Opinion in Plant Biology** 2011, **14**:24–30

This review comes from a themed issue on  
 Growth and development  
 Edited by Jiayang Li and Nam-Hai Chua

Available online 25th September 2010

1369-5266/\$ – see front matter  
 Published by Elsevier Ltd.

DOI [10.1016/j.pbi.2010.08.012](https://doi.org/10.1016/j.pbi.2010.08.012)

### Introduction

Coordination of cell division and differentiation is required for the development of multicellular organisms, which start as single cell zygotes. In plants, organs are continuously generated at the flanks of the meristem, where a population of pluripotent stem cells resides. At the center of the dome-shaped shoot apical meristem (SAM), cells divide slowly to generate more cells that will be used for lateral organs. When stem cells divide, daughter cells are shifted towards the periphery where organ initiation and/or further division occur. Cell fate is therefore connected to a cell's location within the meristematic dome.

During vegetative growth, the SAM continues to produce leaves, which initiate as dorsiventral structures from the flank of the meristem. Leaves capture light energy and use it to convert carbon dioxide and water into sugars and oxygen that will be the energy source for the whole plant. Thus, a leaf needs an optimal structure to maximize light

capture and be efficient in gas exchange. At the same time, leaves have evolved mechanisms, such as dissection, to protect themselves from excessive damage from insects, wind or sun. While leaves are often large and serve as the major photosynthetic structures of plants during vegetative growth, as the plant becomes reproductive, the leaves become small and subtend flowers or branches. The process of leaf development, from initiation to patterning, involves coordinated regulation among transcription factors, small RNAs and hormones. In this review, we will discuss the mechanisms underlying the transition to determinacy during leaf initiation and how polarity is maintained during leaf outgrowth.

### Leaf initiation: a switch to determinacy

The pluripotent state of the SAM is characterized by expression of Class I *KNOTTED1-LIKE HOMEODOMAIN* (*KNOX*) genes [1,2]. Loss-of-function mutants of the *KNOX* genes, *knotted1* (*kn1*) in maize and *SHOOTMERISTEMLESS* (*STM*) in Arabidopsis, show loss of meristem function to various degrees indicating that these genes are required for proper maintenance of the SAM [3,4]. One of the earliest indications of leaf development is the down-regulation of *KNOX* genes at the site of incipient primordia. Through this repression, a switch from an indeterminate to a determinate fate occurs in the small group of cells that will be the immediate precursor of the leaf primordium. *KNOX* transcription factors regulate two phytohormones, gibberellin (GA) and cytokinin. GA promotes cell elongation and differentiation while cytokinin promotes cell proliferation within the meristem. Thus, a high cytokinin to low GA ratio is important to maintain the SAM in an indeterminate state. *KNOX* proteins directly downregulate the *GA20-oxidase* gene, an enzyme encoding a rate-limiting step in GA biosynthesis [5,6]. Inactivation of GA is also achieved through accumulation of the *GA2-oxidase* by *KNOX* proteins [7]. On the other hand, *KNOX* proteins upregulate the expression of cytokinin biosynthesis genes to achieve high levels of cytokinin [8,9]. The downregulation of *KNOX* genes from the incipient primordium will lead to a low cytokinin to high GA ratio promoting the switch from an indeterminate to a determinate state.

Polar auxin transport is considered important for the downregulation of *KNOX* genes in the incipient primordium in both maize and Arabidopsis. Maize apices grown in the presence of an auxin transport inhibitor fail to show *KNOX* downregulation within the meristem and fail to initiate leaves [10]. The *bobber1* mutant of Arabidopsis, which is blocked at the globular stage of development, expresses *STM* throughout the apical half of the embryo

and auxin activity, as visualized by DR5, is not localized [11<sup>•</sup>]. In addition, real-time imaging of developing apices showed localized auxin activity correlates with the site of *KNOX* downregulation [12].

Another pathway involved in *KNOX* repression is mediated by the *ARP* genes, named after *ASYMMETRIC LEAF1* (*AS1*) from *Arabidopsis*, *rough sheath2* (*rs2*) from maize, and *phantastica* (*phan*) from *Antirrhinum*. These genes encode MYB transcription factors and are expressed in lateral organ founder cells where they repress *KNOX* expression (reviewed in [13]). *AS1* forms a repressor complex with the LOB domain protein *AS2* that directly binds to specific regions of *KNOX* promoters [14]. The repressor complex also includes the predicted RNA binding protein *RIK* and the chromatin-remodeling protein *HIRA*, suggesting formation of a repressed chromatin state at the targeted *KNOX* locus during organogenesis [14,15]. In addition, genetic experiments suggest that *AS1* function converges with auxin to repress the *KNOX* gene *BREVIPEDICELLUS* (*BP*) [16]. Thus, multiple levels of regulation and many partners are likely to keep *KNOX* genes out of the leaf initiation site.


### Asymmetric growth of the leaf: which side is up?

Starting as a bulge on the flank of the SAM, the newly initiated leaf becomes asymmetric in several axes: the adaxial–abaxial, medial–lateral, and proximal–distal (Figure 1). The meristem provides leaves with an inherent sidedness, with the adaxial side adjacent to the meristem and the proximal end attached to the

meristem. Polarization along these axes eventually leads to the asymmetric distribution of cell types in the mature leaf that is critical for the physiology of the plant. For example, the adaxial side of the leaf blade often consists of cells specialized for light harvesting while the abaxial side contains cells involved in gas exchange to maximize photosynthesis. Similarly, the distal end of a pea leaf contains the tendrils that cling to support structures and the proximal end of a grass leaf is the sheath, specialized for holding the leaf to the stem.


Characterization of polarity mutants led to the discovery of a complex regulatory network underlying the process of polarity establishment and leaf outgrowth. Key to this discovery was analysis of the dominant mutants, *phavoluta* (*phv*), *revoluta* (*rev*) and *phabulosa* (*phb*) in *Arabidopsis* and *Rolled1* (*Rld1*) in maize [17–19]. The genes defined by these mutants encode class III HOMEODOMAIN LEUCINE ZIPPER (HD-ZIPIII) proteins [18,20,21]. Expression of this gene family is restricted to the adaxial side of the developing leaf and extends into the SAM in a pattern that predicts phyllotaxy [18]. In mutants lacking HD-ZIPIII function, abaxial fate dominates and the SAM fails to form. Gain-of-function mutants occur due to mutations in a miR165/166 binding site within the gene preventing the transcripts from being cleaved [22,23]. In non-mutant leaves, miR165/166 is expressed towards the abaxial side limiting the expression of *HD-ZIPIII* to the adaxial domain of the leaf (Figure 2). In the dominant mutants, *HD-ZIPIII* expression spreads throughout the leaf resulting in adaxialization.

Figure 1



Leaf initiation in (a) *Arabidopsis* and (b) maize. Class I *KNOX* genes are expressed throughout the SAM keeping the cells in an indeterminate state [1,2]. At the site of incipient primordia, downregulation of *KNOX* genes occurs which leads to a switch from an indeterminate to a determinate fate in the small group of cells (founder cells) that will be the immediate precursor of the leaf primordium. The developing leaf becomes asymmetric in several axes: proximal–distal, adaxial–abaxial and medial–lateral. This polarization leads to asymmetric distribution of cell types in the mature leaf. In maize, a leaf develops in a ring-like structure where the proximal end will differentiate into sheath tissue which wraps around the stem. *KNOX* expression domain is colored in grey. P: proximal, D: distal, M: medial, L: lateral, Ad: adaxial, and Ab: abaxial.

Figure 2



Establishing adaxial–abaxial polarity in a developing leaf. *HD-ZIPIII* is expressed in the adaxial domain [18] while *KAN* sets up the abaxial domain of the leaf [24,25]. The antagonistic relationship of these two gene families is established by several layers of regulation. *miR165/166* negatively regulates *HD-ZIPIII* to prevent expression on the abaxial side of the leaf [22,23] while the *AS1/AS2* complex promotes the expression of *HD-ZIPIII* in the adaxial domain [37]. The *AS1/AS2* complex also represses the expression of *miR165/166*, *KAN* and *YAB* [37,38]. The *ARF3/4* transcription factors promote the abaxial fate and the tasiRNAs prevent these genes from being expressed adaxially [47,48\*\*].

Antagonistic to the *HD-ZIPIII* genes are also the *KANADI* (*KAN*) gene family members, expressed on the abaxial side of the leaf primordia [24,25] (Figure 2). Loss-of-function *kan* mutants are adaxialized and *HD-ZIPIII* genes are expressed throughout the leaf whereas *KAN* overexpression causes abaxialization. The *YABBY* (*YAB*) genes function relatively later in leaf development and are considered to act downstream of the *KAN* genes. In Arabidopsis, *YAB* genes expression is abaxial and adaxialization occurs when multiple *yab* mutations are combined [26,27\*].

Three recent reports highlight *kan* loss-of-function mutants in maize and rice. The rice mutant, *shallot-like1* (*sll1*) has rolled leaves due to loss of sclerenchymatous cells on the abaxial sides of the veins [28\*]. Ligules, normally only adaxial, are also found abaxial. The *milkweed pod1* (*mwp1*) mutant of maize affects the sheaths of leaves in patches and not the blade, although the gene is expressed in both sheath and blade [29\*]. The *HD-ZIPIII* gene *Rld1* is misexpressed in these patches. Sheaths are narrower as are leaf-like organs of the flower [30\*]. Prophylls, which are made by the fusion of the first two leaves of the axillary branch, are thread-like and unfused. The narrow leaves in *mwp1* mutants support the model of

Waites and Hudson that proper abaxial–adaxial patterning is required for lamina outgrowth [31].

Genes that were identified as negative regulators of *KNOX* genes are also implicated in polarity. The first polarity mutant characterized was the Antirrhinum *phan*, which has a mutation in a gene orthologous to *AS1* in Arabidopsis [31,32]. Leaves of the *phan* mutant display abaxialization. Mutations in the homologous genes Arabidopsis *AS1* and maize *rs2*, rarely have defects in leaf polarity [33–35]; however, *AS1* plays a role in abaxial–adaxial polarity formation by forming a complex with *AS2* [36–38]. The *AS1/AS2* complex positively regulates *HD-ZIPIII* expression and suppresses *KAN* and *YAB* expression [37] (Figure 2). The positive regulation of *HD-ZIPIII* by the *AS1/AS2* complex is also achieved through reducing *miR165/166* expression [38]. A direct interaction of *AS2* and *KAN* was shown through a novel dominant allele. *as2-5d* has polarity defects that resemble *kan* mutants [39]. *KAN* directly binds to the *AS2* promoter but does not bind to the promoter in the dominant mutant, thereby resulting in high *AS2* levels. The high levels of *AS2* in turn, directly or indirectly, negatively regulate *KAN*. These results may explain why *phan*, which is not expressed in a polar fashion, has a polarity defect. Perhaps the loss of *phan* leads to an increase in *KAN* and thus abaxialization.

Other polarity determinants, the auxin response factors *ARF3/ETTIN* and *ARF4*, were identified as suppressors of a *KAN* overexpressing plant [40]. They function downstream of *KAN* but additional regulation is achieved by trans-acting short interfering RNAs (tasiRNAs) [41,42]. TasiRNAs are generated from *TAS* genes, which do not encode a protein but function as an RNA. The production of tasiRNAs requires a number of proteins including *AGO7*, *SGS3*, *RDR6* and *DCL4* [43–46]. *ARF3* and *ARF4* mRNAs are cleaved and degraded by these tasiRNAs [47]. tasiRNAs targeting the ARFs are produced at the adaxial side of the primordium and prevent ARF activity in the adaxial domain [48\*\*] (Figure 2). The two small RNAs, tasiRNA and *miR165/166*, show opposite polar distribution in leaf primordia and are thought to establish the adaxial–abaxial axis in leaf development [42].

Mutations in the tasiRNA biogenesis pathway or plants with *TAS3*-insensitive *ARF3* do not show striking polarity defects in Arabidopsis [47,49], but they do in the grasses. In rice, mutants in *RDR6* (*shootless2*), *AGO7* (*shootless4*) and *DCL4* (*shoot organization2*) produce shootless embryos or filamentous leaves [50,51]. In maize, the *leafbladeless1* (*lbl1*) mutant, which carries a mutation in *SGS3*, shows radialized abaxialization [42]. The maize *ago7* mutant, which was recently shown to be *ragged seedling2* (*rgd2*) [52\*\*], has cylindrical leaves, but surprisingly, no loss of dorsiventrality [53]. *rgd2-r* mutants have less tasiARF and

an increase in *ARF3A*, but expression of *ARF3A* is still polarized in both *rgd2* and in *lbl1* mutants [52<sup>••</sup>]. The difference in polarity between *lbl1* and *rgd2* mutants suggests that LBL1 does more than accumulate tasiRNAs. The regulation of leaf polarity is likely through *AGO1*, which is increased in *lbl1* but not *rgd2* mutants [52<sup>••</sup>] and accumulates adaxially in maize [54].

### The role of *KNOX* genes in proximal distal polarity and leaf complexity


Just as the absence of *KNOX* genes plays a role in defining the position of a leaf within the meristem, expression of *KNOX* genes plays a critical role in elaborating leaf shape in many species (reviewed in [55,56]). The timing and location of *KNOX* expression in the leaf turns out to be critical for the outcome.

Maize leaves have a proximal sheath that wraps tightly around the stem and a distal blade that lies back to optimize photosynthesis. At the junction of the sheath and blade is a ligule region that contains an adaxial ligule fringe and two auricles (Figure 1). Maize, like other plants with simple leaves, does not express *KNOX* genes in the leaf; however, a number of dominant mutants exist in which *KNOX* genes are expressed in the leaf [57,58]. For most of these mutants, *Liguleless3*, *Roughsheath1*, *Gnarley*, and some *Kn1* alleles, the ligule region is displaced to a more distal position. In cases where this has been studied by *in situ*, *KNOX* expression extends into the sheath from the adjoining stem [59,60]. In some *Kn1* alleles, however, the misexpression is not contiguous with expression in the meristem, but is in the veins of the blade or along the margins. The outcomes are quite different depending on when and where expression occurs (Figure 3). When *kn1* is expressed in the lateral veins, the cells adopt sheath and auricle fates and form protrusions known as knots [61]. However, when *kn1* is expressed at the margin, flaps of tissue grow out from the margin that are sheath and auricle in leaf cell type. When *kn1* is expressed at the very tip of the growing leaf, growth is arrested and the leaf becomes bifurcated [62<sup>•</sup>]. These phenotypes suggest that *kn1* regulates proximal distal patterning [62<sup>•</sup>].

Another mutant with defects in proximal distal identity is *blade on petiole (bop)* of Arabidopsis. Ectopic blade tissue is found in a proximal position, along the petiole (Figure 3) [63,64]. *bop* mutants accumulate *KNOX* genes in the leaf and the BOP1/BOP2 complex directly and positively regulates AS2 [65<sup>•</sup>]. Thus, the absence of BOP function leads to less AS2 and more *KNOX*, yet the phenotype, a proximal (petiole) to distal (blade) transformation, is opposite to that of the maize mutants [57].

Tomato leaves have a terminal leaflet and alternating leaflets from the main rachis. Second order leaflets occur from the first order leaflets, but not from the terminal leaflet. *KNOX* genes are expressed in leaves and

Figure 3



Ectopic *KNOX* expression in leaves leads to a different outcome depending on the location and timing. In normal maize, *kn1* expression is excluded from the developing leaf. The *Kn1-DL* and the *Kn1-O* alleles are shown here which have ectopic *kn1* expression in the leaves [61,62<sup>•</sup>]. In the *Kn1-DL* allele, *kn1* is expressed at the very tip of the growing leaf leading to arrest in growth and the leaf bifurcation (b). In the *Kn1-O* alleles, the ectopic *kn1* expression results in knots (shown with an arrow) which consists of cells with sheath and auricle fates (c). *KNOX* expression is also excluded from the developing leaf of Arabidopsis. When *KNOX* genes are turned on in leaves, as in the case of *bop* mutants, ectopic blade tissue appears at a proximal position, along the petioles (indicated with an arrow) (d,e) [63,64].

additional expression often leads to further dissection of the first order leaflets (reviewed in [56]). Shani *et al.* expressed the STM ortholog of tomato, *TKn2*, behind the *FIL* promoter, which gives expression in young leaf primordia but not the meristem [66]. Constructs either included or did not include an *EAR* repressor domain [67<sup>••</sup>]. Surprisingly, both constructs led to simpler leaves, but for different reasons. Plants that expressed *FIL:TKn2-EAR* prematurely differentiated, before developing leaflets, whereas expression of *FIL:TKn2* kept leaves in a perpetual early plastochron state, unable to expand the lamina. They also used promoters that function at different times of leaf maturation. Only promoters that function during the primary morphogenesis state increased the complexity of the leaf [67<sup>••</sup>].

One of the facilitators of leaf diversity generated by *KNOX* is likely to be the *CUP-SHAPED COTYLEDON (CUC)* gene family, which is expressed at the boundary of leaves and leaflets [68,69]. Dex induction of a *KN1-GR* fusion led to an increase in lobing and an increase in *CUC* gene expression in *Cardamine* [70]. Loss of *CUC* gene function led to a less dissected leaf in a number of species including pea, potato, tomato, *Cardamine* and *Aquilegia*



[70]. However, overexpression of a microRNA resistant version of the tomato *CUC* gene, *GOBLET* also had fewer leaflets, presumably because of leaflet fusion [71]. Clearly the context of both *KNOX* and *CUC* expression is critical for the genesis of leaf shape, whether they make a leaf dissected or simple, and whether they displace proximal or distal tissues.

## Conclusion

Diverse species often utilize the same set of regulators to elaborate structures. Slight changes in timing and expression pattern of these regulators are likely to result in dramatic morphological variation. Plant scientists are in position to take advantage of the knowledge gained from model organisms to begin to probe evolutionary change, and to move from the transcription factor regulators to understanding the cellular processes of growth.

## Acknowledgements

Funding was provided by CSREES (2008-01082). We wish to thank Toshi Foster for helpful comments.

## References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Smith LG, Greene B, Veit B, Hake S: **A dominant mutation in the maize homeobox gene, *Knotted-1*, causes its ectopic expression in leaf cells with altered fates.** *Development* 1992, **116**:21-30.
  2. Jackson D, Veit B, Hake S: **Expression of maize *KNOTTED1* related homeobox genes in the shoot apical meristem predicts patterns of morphogenesis in the vegetative shoot.** *Development* 1994, **120**:405-413.
  3. Long JA, Moan EI, Medford JI, Barton MK: **A member of the *KNOTTED* class of homeodomain proteins encoded by the *SHOOTMERISTEMLESS* gene of *Arabidopsis*.** *Nature* 1996, **379**:66-69.
  4. Vollbrecht E, Reiser L, Hake S: **Shoot meristem size is dependent on inbred background and presence of the maize homeobox gene, *knotted1*.** *Development* 2000, **127**:3161-3172.
  5. Sakamoto T, Kamiya N, Ueguchi-Tanaka M, Iwahori S, Matsuoka M: ***KNOX* homeodomain protein directly suppresses the expression of a gibberellin biosynthetic gene in the tobacco shoot apical meristem.** *Genes Dev* 2001, **15**:581-590.
  6. Hay A, Kaur H, Phillips A, Hedden P, Hake S, Tsiantis M: **The gibberellin pathway mediates *KNOTTED1*-type homeobox function in plants with different body plans.** *Curr Biol* 2002, **12**:1557-1565.
  7. Bolduc N, Hake S: **The maize transcription factor *KNOTTED1* directly regulates the gibberellin catabolism gene *ga2ox1*.** *Plant Cell* 2009, **21**:1647-1658.
- This work shows the direct activation of *ga2ox1* by KN1 in maize, suggesting that KN1 negatively modulates the accumulation of GA by activating this catabolism enzyme.
8. Jasinski S, Piazza P, Craft J, Hay A, Woolley L, Rieu I, Phillips A, Hedden P, Tsiantis M: ***KNOX* action in *Arabidopsis* is mediated by coordinate regulation of cytokinin and gibberellin activities.** *Curr Biol* 2005, **15**:1560-1565.
  9. Yanai O, Shani E, Dolezal K, Tarkowski P, Sablowski R, Sandberg G, Samach A, Ori N: ***Arabidopsis KNOX1* proteins activate cytokinin biosynthesis.** *Curr Biol* 2005, **15**:1566-1571.
  10. Scanlon MJ: **The polar auxin transport inhibitor N-1-naphthylphthalamic acid disrupts leaf initiation, *KNOX* protein regulation, and formation of leaf margins in maize.** *Plant Physiol* 2003, **133**:597-605.
  11. Jurkuta RJ, Kaplinsky NJ, Spindel JE, Barton MK: **Partitioning the apical domain of the *Arabidopsis* embryo requires the *BOBBER1* NudC domain protein.** *Plant Cell* 2009, **21**:1957-1971.
- In this paper, the authors identified the *bob1* mutation which arrests development at the globular stage. The mutants show *STM* expression at the apical half of the embryo but fail to express *ANT* and local maxima of DR5 is not observed.
12. Heisler MG, Ohno C, Das P, Sieber P, Reddy GV, Long JA, Meyerowitz EM: **Patterns of auxin transport and gene expression during primordium development revealed by live imaging of the *Arabidopsis* inflorescence meristem.** *Curr Biol* 2005, **15**:1899-1911.
  13. Byrne ME: **Shoot development—genetic interactions in the meristem.** *Biochem Soc Trans* 2005, **33**:1499-1501.
  14. Guo M, Thomas J, Collins G, Timmermans MC: **Direct repression of *KNOX* loci by the *ASYMMETRIC LEAVES1* complex of *Arabidopsis*.** *Plant Cell* 2008, **20**:48-58.
  15. Phelps-Durr TL, Thomas J, Vahab P, Timmermans MC: **Maize rough sheath2 and its *Arabidopsis* orthologue *ASYMMETRIC LEAVES1* interact with HIRA, a predicted histone chaperone, to maintain *knox* gene silencing and determinacy during organogenesis.** *Plant Cell* 2005, **17**:2886-2898.
  16. Hay A, Barkoulas M, Tsiantis M: ***ASYMMETRIC LEAVES1* and auxin activities converge to repress *BREVIPEDICELLUS* expression and promote leaf development in *Arabidopsis*.** *Development* 2006, **133**:3955-3961.
  17. Otsuga D, DeGuzman B, Prigge MJ, Drews GN, Clark SE: ***REVOLUTA* regulates meristem initiation at lateral positions.** *Plant J* 2001, **25**:223-236.
  18. McConnell JR, Emery JF, Eshed Y, Bao N, Bowman J, Barton MK: **Role of *PHABULOSA* and *PHAVOLUTA* in determining radial patterning in shoots.** *Nature* 2001, **411**:709-713.
  19. Nelson JM, Lane B, Freeling M: **Expression of a mutant maize gene in the ventral leaf epidermis is sufficient to signal a switch of the leaf's dorsoventral axis.** *Development* 2002, **129**:4581-4589.
  20. Emery JF, Floyd SK, Alvarez J, Eshed Y, Hawker NP, Izhaki A, Baum SF, Bowman JL: **Radial patterning of *Arabidopsis* shoots by Class III HD-ZIP genes is mediated by miRNA regulation.** *Curr Biol* 2003, **13**:1768-1774.
  21. Juarez MT, Kui JS, Thomas J, Heller BA, Timmermans MC: **microRNA-mediated repression of *rolled leaf1* specifies maize leaf polarity.** *Nature* 2004, **428**:84-88.
  22. Rhoades MW, Reinhart BJ, Lim LP, Burge CB, Bartel B, Bartel DP: **Prediction of plant microRNA targets.** *Cell* 2002, **110**:513-520.
  23. Mallory AC, Reinhart BJ, Jones-Rhoades M, Tang G, Zamore PD, Barton MK, Bartel DP: **MicroRNA control of *PHABULOSA* in leaf development: importance of pairing to the microRNA 5' region.** *EMBO J* 2004, **23**:3356-3364.
  24. Kerstetter RA, Bollman K, Taylor RA, Bombliks K, Poethig RS: ***KANADI* regulates organ polarity in *Arabidopsis*.** *Nature* 2001, **411**:706-709.
  25. Eshed Y, Baum SF, Perea JV, Bowman JL: **Establishment of polarity in lateral organs of plants.** *Curr Biol* 2001, **11**:1251-1260.
  26. Eshed Y, Izhaki A, Baum SF, Floyd SK, Bowman JL: **Asymmetric leaf development and blade expansion in *Arabidopsis* are mediated by *KANADI* and *YABBY* activities.** *Development* 2004, **131**:2997-3006.
  27. Stähle MI, Kuehlich J, Staron L, von Arnim AG, Golz JF: ***YABBYs* and the transcriptional corepressors *LEUNIG* and *LEUNIG\_HOMOLOG* maintain leaf polarity and meristem activity in *Arabidopsis*.** *Plant Cell* 2009, **21**:3105-3118.
- This work shows that YAB proteins physically interact with components of a transcriptional repressor complex that include LUG, LUH, SEUSS, and

related SEUSS-LIKE proteins. Genetic analysis suggests that LUG-YAB complex promotes adaxial cell identity in leaves as well as SAM initiation and maintenance.

28. Zhang GH, Xu Q, Zhu XD, Qian Q, Xue HW: **SHALLOT-LIKE1 is a KANADI transcription factor that modulates rice leaf rolling by regulating leaf abaxial cell development.** *Plant Cell* 2009, **21**:719-735.

The authors isolated and functionally characterized the *SLL1* gene, which encodes a KAN family member. Abaxial accumulation of the transcript and deficient abaxial development in the mutant suggests that *SLL1* controls proper abaxial development, which is important in modulating leaf rolling.

29. Candela H, Johnston R, Gerhold A, Foster T, Hake S: **The milkweed pod1 gene encodes a KANADI protein that is required for abaxial-adaxial patterning in maize leaves.** *Plant Cell* 2008, **20**:2073-2087.

In this paper, the authors describe the *mwp1* mutant in maize which shows sectors of adaxialization in the sheath. The *mwp1* gene encodes a KAN transcription factor and the transcript is found in the abaxial domain of the leaf.

30. Johnston R, Candela H, Hake S, Foster T: **The maize milkweed pod1 mutant reveals a mechanism to modify organ morphology.** *Genesis* 2010, **48**:416-423.

This paper extends the characterization of the *mwp1* mutant [29] to various organs including prophyll and silks. The authors show that fused organ development and lateral growth are dependent on adaxial-abaxial polarity.

31. Waites R, Hudson A: **phantastica: a gene required for dorsoventrality of leaves in *Antirrhinum majus*.** *Development* 1995, **121**:2143-2154.
32. Waites R, Selvadurai HR, Oliver IR, Hudson A: **The PHANTASTICA gene encodes a MYB transcription factor involved in growth and dorsoventrality of lateral organs in *Antirrhinum*.** *Cell* 1998, **93**:779-789.
33. Timmermans MC, Hudson A, Becraft PW, Nelson T: **ROUGH SHEATH2: a Myb protein that represses knox homeobox genes in maize lateral organ primordia.** *Science* 1999, **284**:151-153.
34. Tsiantis M, Schneeberger R, Golz JF, Freeling M, Langdale JA: **The maize rough sheath2 gene and leaf development programs in monocot and dicot plants.** *Science* 1999, **284**:154-156.
35. Byrne ME, Barley R, Curtis M, Arroyo JM, Dunham M, Hudson A, Martienssen RA: **asymmetric leaves 1 mediates leaf patterning and stem cell function in Arabidopsis.** *Nature* 2000, **408**:967-971.
36. Lin WC, Shuai B, Springer PS: **The Arabidopsis LATERAL ORGAN BOUNDARIES-domain gene ASYMMETRIC LEAVES2 functions in the repression of KNOX gene expression and in adaxial-abaxial patterning.** *Plant Cell* 2003, **15**:2241-2252.
37. Fu Y, Xu L, Xu B, Yang L, Ling Q, Wang H, Huang H: **Genetic interactions between leaf polarity-controlling genes and ASYMMETRIC LEAVES1 and 2 in Arabidopsis leaf patterning.** *Plant Cell Physiol* 2007, **48**:724-735.
38. Li H, Xu L, Wang H, Yuan Z, Cao X, Yang Z, Zhang D, Xu Y, Huang H: **The Putative RNA-dependent RNA polymerase RDR6 acts synergistically with ASYMMETRIC LEAVES1 and 2 to repress BREVIPEDICELLUS and MicroRNA165/166 in Arabidopsis leaf development.** *Plant Cell* 2005, **17**:2157-2171.
39. Wu G, Lin WC, Huang T, Poethig RS, Springer PS, Kerstetter RA: **KANADI1 regulates adaxial-abaxial polarity in Arabidopsis by directly repressing the transcription of ASYMMETRIC LEAVES2.** *Proc Natl Acad Sci USA* 2008, **105**:16392-16397.
40. Pekker I, Alvarez JP, Eshed Y: **Auxin response factors mediate Arabidopsis organ asymmetry via modulation of KANADI activity.** *Plant Cell* 2005, **17**:2899-2910.
41. Garcia D, Collier SA, Byrne ME, Martienssen RA: **Specification of leaf polarity in Arabidopsis via the trans-acting siRNA pathway.** *Curr Biol* 2006, **16**:933-938.
42. Nogueira FT, Madi S, Chitwood DH, Juarez MT, C TM: **Two small regulatory RNAs establish opposing fates of a developmental axis.** *Genes Dev* 2007, **21**:750-755.

43. Peragine A, Yoshikawa M, Wu G, Albrecht HL, Poethig RS: **SGS3 and SGS2/SDE1/RDR6 are required for juvenile development and the production of trans-acting siRNAs in Arabidopsis.** *Genes Dev* 2004, **18**:2368-2379.
  44. Allen E, Xie Z, Gustafson A, Carrington J: **micro-directed phasing during trans-acting siRNA biogenesis in plants.** *Cell* 2005, **121**:207-221.
  45. Yoshikawa M, Peragine A, Park MY, Poethig RS: **A pathway for the biogenesis of trans-acting siRNAs in Arabidopsis.** *Genes Dev* 2005, **19**:164-175.
  46. Adenot X, Elmayan T, Lauressergues D, Boutet S, Bouché N, Gascioli V, Vaucheret H: **DRB4-dependent TAS3 trans-acting siRNAs control leaf morphology through AGO7.** *Curr Biol* 2006, **16**:927-932.
  47. Fahlgren N, Montgomery TA, Howell MD, Allen E, Dvorak SK, Alexander AL, Carrington JC: **Regulation of AUXIN RESPONSE FACTOR3 by TAS3 ta-siRNA affects developmental timing and patterning in Arabidopsis.** *Curr Biol* 2006, **16**:939-944.
  48. Chitwood DH, Nogueira FT, Howell MD, Montgomery TA, Carrington JC, Timmermans MC: **Pattern formation via small RNA mobility.** *Genes Dev* 2009, **23**:549-554.
- This work suggests that *tasiARFs* move from the adaxial side of the leaf to the abaxial side, which results in maintaining the polarized expression of *ARF3* to the abaxial side of the leaf.
49. Hunter C, Willmann MR, Wu G, Yoshikawa M, de la Luz Gutierrez-Nava M, Poethig RS: **Trans-acting siRNA-mediated repression of ETTIN and ARF4 regulates heteroblasty in Arabidopsis.** *Development* 2006, **133**:2973-2981.
  50. Nagasaki H, Itoh J, Hayashi K, Hibara K, Satoh-Nagasawa N, Nosaka M, Mukouhata M, Ashikari M, Kitano H, Matsuoka M *et al.*: **The small interfering RNA production pathway is required for shoot meristem initiation in rice.** *Proc Natl Acad Sci USA* 2007, **104**:14867-14871.
  51. Itoh J, Sato Y, Nagato Y: **The SHOOT ORGANIZATION2 gene coordinates leaf domain development along the central-marginal axis in rice.** *Plant Cell Physiol* 2008, **49**:1226-1236.
  52. Douglas RN, Wiley D, Sarkar A, Springer N, Timmermans MC, Scanlon MJ: **Ragged seedling2 encodes an ARGONAUTE7-like protein required for mediolateral expansion, but not dorsiventrality, of maize leaves.** *Plant Cell* 2010, **22**:1441-1451.
- The authors show that *ragged seedling2* is a mutation in the maize *ago7* gene. The leaves are radial, but still have adaxial-abaxial polarity and *arf3a* expression is still polarized suggesting that *tasiARF* is not required for the polarization of *arf3a*. They also show that maize *ago1* transcripts overaccumulate in *lb1* but not in *rgd2* mutants, suggesting that upregulation of *ago1* contributes to abaxialized leaf formation in *lb1*.
53. Henderson DC, Muehlbauer GJ, Scanlon MJ: **Radial leaves of the maize mutant ragged seedling2 retain dorsiventral anatomy.** *Dev Biol* 2005, **282**:455-466.
  54. Brooks LR, Strable J, Zhang X, Ohtsuka K, Zhou R, Sarkar A, Hargreaves S, Elshire RJ, Eudy D, Pawlowska T *et al.*: **Microdissection of shoot meristem functional domains.** *PLoS Genet* 2009, **5**:e1000476.
  55. Blein T, Hasson A, Laufs P: **Leaf development: what it needs to be complex.** *Curr Opin Plant Biol* 2010, **13**:75-82.
  56. Efroni I, Eshed Y, Lifschitz E: **Morphogenesis of simple and compound leaves: a critical review.** *Plant Cell* 2010, **22**:1019-1032.
  57. Freeling M: **A conceptual framework for maize leaf development.** *Dev Biol* 1992, **153**:44-58.
  58. Hake S, Smith HMS, Holtan H, Magnani E, Mele G, Ramirez J: **The role of KNOX genes in plant development.** *Annu Rev Cell Dev Biol* 2004, **20**:125-151.
  59. Schneeberger RG, Becraft PW, Hake S, Freeling M: **Ectopic expression of the knox homeo box gene rough sheath1 alters cell fate in the maize leaf.** *Genes Dev* 1995, **9**:2292-2304.
  60. Foster T, Yamaguchi J, Wong BC, Veit B, Hake S: **Gnarley is a dominant mutation in the knox4 homeobox gene affecting cell shape and identity.** *Plant Cell* 1999, **11**:1239-1252.

61. Sinha N, Hake S: **The *Knotted* leaf blade is a mosaic of blade, sheath, and auricle identities.** *Dev Genet* 1994, **15**:401-414.
62. Ramirez J, Bolduc N, Lisch D, Hake S: **Distal expression of *knotted1* in maize leaves leads to re-establishment of proximal/distal patterning and leaf dissection.** *Plant Physiol* 2009, **151**:1878-1888.  
 The authors describe a new dominant allele of maize *kn1*, *Kn1-DL*, which contains a transposon insertion in the promoter in addition to a tandem duplication of the *kn1* locus. When *kn1* is misexpressed along the margins, flaps of sheath-like tissue form along the margins while the expression in the distal tip leads to premature growth and leaf bifurcates. The gain-of-function phenotypes suggest that *kn1* establishes proximal/distal patterning.
63. Ha CM, Jun JH, Nam HG, Fletcher JC: ***BLADE-ON-PETIOLE1* encodes a BTB/POZ domain protein required for leaf morphogenesis in *Arabidopsis thaliana*.** *Plant Cell Physiol* 2004, **45**:1361-1370.
64. Norberg M, Holmlund M, Nilsson O: **The *BLADE ON PETIOLE* genes act redundantly to control the growth and development of lateral organs.** *Development* 2005, **132**:2203-2213.
65. Jun JH, Ha CM, Fletcher JC: ***BLADE-ON-PETIOLE1* coordinates organ determinacy and axial polarity in arabidopsis by directly activating *ASYMMETRIC LEAVES2*.** *Plant Cell* 2010, **22**:62-76.  
 In this paper, authors show that BOP1 and BOP2 dimerize and function as transcriptional activators. They also show that the *AS2* gene is directly activated by BOP at the base of the leaf, preventing the expression of *KNOX* genes in this region.
66. Lifschitz E, Eviatar T, Rozman A, Shalit A, Goldshmidt A, Amsellem Z, Alvarez JP, Eshed Y: **The tomato *FT* ortholog triggers systemic signals that regulate growth and flowering and substitute for diverse environmental stimuli.** *Proc Natl Acad Sci USA* 2006, **103**:6398-6403.
67. Shani E, Burko Y, Ben-Yaakov L, Berger Y, Amsellem Z, Goldshmidt A, Sharon E, Ori N: **Stage-specific regulation of *Solanum lycopersicum* leaf maturation by class 1 *KNOTTED1*-LIKE *HOMEBOX* proteins.** *Plant Cell* 2009, **21**:3078-3092.  
 The authors dissected the function of *KNOX1* in compound leaf development by expressing the *Tkn2* with and without a repressor domain. Both cases resulted in narrow and simple leaves but due to different reasons. In the presence of the repressor domain, differentiation was accelerated, while leaf development was blocked at the initiation stage without the repressor domain.
68. Souer E, Vanhouwelingen A, Kloos D, Mol J, Koes R: **The no apical meristem gene of petunia is required for pattern formation in embryos and flowers and is expressed at meristem and primordia boundaries.** *Cell* 1996, **85**:159-170.
69. Aida M, Ishida T, Fukaki H, Fujisawa H, Tasaka M: **Genes involved in organ separation in arabidopsis: an analysis of the cup-shaped cotyledon mutant.** *Plant Cell* 1997, **9**:841-857.
70. Blein T, Pulido A, Vialette-Guiraud A, Nikovics K, Morin H, Hay A, Johansen IE, Tsiantis M, Laufs P: **A conserved molecular framework for compound leaf development.** *Science* 2008, **322**:1835-1839.
71. Berger Y, Harpaz-Saad S, Brand A, Melnik H, Sirding N, Alvarez JP, Zinder M, Samach A, Eshed Y, Ori N: **The *NAC*-domain transcription factor *GOBLET* specifies leaflet boundaries in compound tomato leaves.** *Development* 2009, **136**:823-832.